

KPHITE® 7LP

SYSTEMIC FUNGICIDE BACTERICIDE



Product Information:

A systemic fungicide bactericide for the control of downy mildew, phytophthora, pythium, and various other diseases on greenhouse crops, indoor and outdoor ornamentals, bedding plants and forestry applications, and domestic trees including conifers, pines, oaks and turf.

Active Ingredients

ACTIVE INGREDIENTS:

Mono- and dipotassium salts of Phosphorous Acid*
.....56%

OTHER INGREDIENTS:.....44%

TOTAL:.....100%

*Contains 7.03 lbs./gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 4.41 lbs. Phosphorous Acid/gallon.

* **K-PHITE® 7LP's** unique manufacturing process and molecule are protected by the following patents:

Reactor: 4724132

Molecule: 7887616, 8221516, 8088191

MANUFACTURED BY:

Plant Food Systems, Inc.
2827 Union St. P.O. Box 775
Zellwood, FL 32798
800.343.7775

EPA Reg. No. 73806-1
EPA Est. No. 73806-FL-001

NET CONTENTS:

- 2.5 Gallons
- 5 Gallons
- 30 Gallons
- 55 Gallons

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PlantFoodSystems.com

See inside booklet for First Aid and
Additional Precautionary Statements.



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENTS

- If in eyes
- Hold eye open and rinse slowly and gently with water for 15-20 minutes
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should remove PPE/clothing immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

(For containers with contents greater than 5 gallons Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

GENERAL APPLICATION INSTRUCTIONS

Apply K-PHITE 7LP by various application methods, including foliar sprays, trunk injection, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply K-PHITE 7LP with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 of the required volume of water.
2. Add K-PHITE 7LP slowly to the tank and agitate by hydraulic or mechanical means.
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

For 1-gallon (3.785 liters) container sizes: this package can treat up to 34,848 sq. ft. at the maximum treatment rate.

For 2 ½-gallon (9.463 liters) container sizes: this package can treat up to 2 acres at the maximum treatment rate.

For 5-gallon (18.925 liters) container sizes: this package can treat up to 4 acres at the maximum treatment rate.

LOW VOLUME AND ULTRA-LOW VOLUME APPLICATIONS

Label rates are written for conventional application equipment. For LV/ULV equipment, use per acre rate of K-PHITE 7LP and adjust amount of water to equipment needs. Always conduct a phytotoxicity test on a small area before applying to a large area in order to assess any potential risk to plants.

SEED TREATMENT APPLICATION INSTRUCTIONS

K-PHITE 7LP is a water-based liquid formulation designed for easy use as seed treatment, and should not form sediment during normal storage. It may be applied using commercial seed treatment equipment or through the use of a hopper-box, planter-box, slurry-box or other seed treatment equipment at or immediately before planting. Equipment should be calibrated to provide a good mixing of seed and treatment products and provide even distribution and coverage of the seed. Uniform application to seed will give optimum performance. The seed should be sound, well-conditioned, and free from dust and chaff.

K-PHITE 7LP can be pre-diluted with clean water sufficient for coverage and other seed amendments can also be mixed with it as required, and the total volume of slurry adjusted according to the application equipment and seed type. Apply gentle agitation during mixing and use, and do not store the diluted product overnight.

This product may be applied only on seeds for crops listed on this label. Do not use treated seed for food or feed purposes or process for oil.

If this product is intended for commercial seed treatment, treated seed must be labeled in accordance with the requirements of the Federal Seed Act and applicable State seed laws. An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed, or oil purposes. If this product is intended for treatment at planting, treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. To dispose of excess treated seed bury away from streams and bodies of water. A dye is not required if excess seed is disposed of by burial.

SEED TREATMENT APPLICATION RATES

Apply K-PHITE 7LP uniformly to seed at rates of 2-12 fl oz/100 lb of seed. Adjust rate of use depending on disease pressure, seed size, density, and ability to absorb liquid.

USE RESTRICTIONS AND PRECAUTIONS

- Store treated seed in cool dry conditions
- Any treated seed which is stored until the following season must be re-tested for germination and vigor prior to planting.

K-PHITE 7LP is compatible with most commonly used seed amendment products, and it is advisable not to tank-mix with any other product unless adequate physical compatibility and phytotoxicity (germination) tests have been conducted and proven to be satisfactory. See compatibility statement below. Consult your supplier or advisor for further information.

COMPATIBILITY

Mixing K-PHITE 7LP with certain surfactants, foliar fertilizers or other pesticides may cause crop injury. K-PHITE 7LP is a slightly acidic buffer solution. Avoid mixing K-PHITE 7LP with strongly acidic or alkaline materials. Do not tank mix without first testing the mixture's compatibility nor apply it without assessing its safety to the crop (phytotoxicity). It is not possible to test all material combinations and environmental tank mix combinations. Always conduct a test with the intended tank mix on a small area prior to any large-scale applications. To determine if a combination is phytotoxic to a specific crop, spray a small area of foliage and fruit, and then evaluate 3-7 days later for visual symptoms.

Test the compatibility of spray adjuvants (i.e. stickers, spreaders, wetting agents) with K-PHITE 7LP before use. Do not use acidifying type compatibility agents.

To determine the compatibility of K-PHITE 7LP with other products, use a jar compatibility test. Add the correct proportions of each product and the appropriate quantity of water to clean container, thoroughly mix, then let stand for 3-5 minutes. If the mixture remains in solution or can be remixed readily, the products are considered compatible.

CHEMIGATION APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

Apply this product through microsprinkler, drip, solid set and center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

PREPARATION OF INJECTION EQUIPMENT: Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: Fill tank with 1/2 to 3/4 of desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of K-PHITE 7LP, then the remaining volume of water. Set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the solution of K-PHITE 7LP into the irrigation water line so as to deliver the desired rate per acre. Inject the K-PHITE 7LP solution with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. For other questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: For Microsprinkler and Drip Irrigation Systems: Avoid further irrigation of the treated area for 24 to 48 hours after treatment with K-PHITE 7LP has been completed. For Solid Set and Center Pivot Irrigation Systems: Avoid further irrigation of the K-PHITE 7LP treated area until after the foliage has completely dried. This will help to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure uniformity solution. Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled or drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of chemigation water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut down and adjust the system as needed.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

ORNAMENTALS

K-PHITE 7LP fungicide is used for control of Pythium, Downy Mildew, Powdery Mildew, Phytophthora and various other diseases of ORNAMENTALS grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. Apply K-PHITE 7LP to all ornamental and landscape plants including (but not limited to): Aglaonema, Aphelandra, Arborvitae, Ash, Azaleas, Bougainvilles, Boxwood, Bushes, Cattleya skinneri, Cedars, Ceanothus, Cotoneaster, Cissus, Dffenbachia, Dogwood, Ferns, Ficus, Flowering Trees, Hibiscus, Hard Wood Trees, Holly, Home Lawns, Juniper, Magnolia Trees, Maple, Pines, Oaks, Palms, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Red woods, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Shrubs, Spathiphyllum, Sycamore, Taxus media, Zygocactus.

APPLICATION GUIDELINES:

Disease Prevention Program*: Apply lower rate at 7-28 day intervals. Do not apply K-PHITE 7LP at intervals less than 3 days.

Disease Control Program: Apply higher rate at 7-14 day intervals until control is reached. Under severe circumstances, application can be made at intervals of up to every three days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity.

Do not apply K-PHITE 7LP at intervals less than 3 days.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

DISEASE	APPLICATION METHOD	RATE/APPLICATION
Anthracnose Fire Blight <i>(Erwinia amylovora)</i> Fusarium Geranium Wilt <i>(Ralstonia spp.)</i> Phytophthora Pythium Rhizoctonia Xanthomonas spp <i>(Including campestris)</i>	Foliar Spray	1-2 quarts in a minimum of 100 gallons of water/acre or 1/3 – 2/3 fluid ounce/gallon of water.
	Soil Drench	12-24 fluid ounce/100 gallon of water or 1/8 – 1/4 fluid ounce per gallon of water
	Soil Incorporation	1-2 pints/cubic yard of soil mix. Mix 1 to 2 pints of K-PHITE 7LP with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development make additional applications as a foliar spray or soil drench.
	Irrigation	Overhead: 1-2 quarts/100 gallons of water. Drip: 1-2 quarts/100 gallons of water. Apply with normal irrigation schedule.
	Trunk or Bark Injection	3 teaspoons/linear yard of canopy width or 2 inches of trunk diameter at breast height (DBH). Inject directly into tree. Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Root Dip	2 quarts/100 gallons of water or 2/3 fluid ounce/gallon of water. Dip transplants into diluted solution of K-PHITE 7LP. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
Downy Mildew Powdery Mildew	Foliar Spray	1-2 quarts/100 gallons of water or 1/3 - 2/3 fluid ounce/ gallon of water

* Check with your local Extension Agent if you are unsure these conditions exist.

BEDDING PLANTS

K-PHITE 7LP fungicide is used for control of Downy Mildew, Powdery Mildew, Pythium, Phytophthora and various other diseases of bedding plants grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. Apply K-PHITE 7LP to all bedding plants including: Ageratum, Algerian Ivy, Alyssum, Anthurium, Artemisia, Aster, Baby's Breath, Begonia, Caladium, Carnation, Chrysanthemum, Coleus, Columbine, Daisy, Delphinium, Easter Lily, English Ivy, Foxglove, Gaillardia, Geranium, Gloxinia, Impatiens, Marigold, Pansy, Petunia, Phlox, Pinks, Poinsettia, Primrose, Prostrate Rosemary, Salvia, Snapdragon, Verbena, Vinca, Zinnia, etc.

APPLICATION GUIDELINES:

Disease Prevention Program*: Apply lower rate at 7-28 day intervals. Do not apply K-PHITE 7LP at intervals less than 3 days.

Disease Control Program: Apply higher rate at 7-14 day intervals until control is reached. Under severe circumstances, application can be made at intervals of up to every three days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity.

Do not apply K-PHITE 7LP at intervals less than 3 days.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

DISEASE	APPLICATION METHOD	RATE/APPLICATION
Anthracnose Fire Blight (<i>Erwinia amylovora</i>) Fusarium Geranium Wilt (<i>Ralstonia spp.</i>) Phytophthora Pythium Rhizoctonia	Foliar Spray	1-2 quarts in a minimum of 100 gallons of water/acre or 1/3 – 2/3 fluid ounce/gallon of water. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.
	Root Dip	2 quarts/100 gallons of water or 2/3 fluid ounces/gallon of water. Dip transplants into diluted solution of K-PHITE 7LP. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
	Soil Drench	12-24 fluid ounces/100 gallons of water of 1/8 to 1/4 fluid ounce/gallon of water.
	Irrigation	Overhead: 1-2 quarts/100 gallons of water. Drip: 1-2 quarts/100 gallons of water. Apply with normal irrigation schedule.
Downy Mildew Powdery Mildew	Foliar Spray	1-2 quarts/100 gallons water or 1/3 - 2/3 fluid ounce/gallon of water. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year.

* Check with your local Extension Agent if you are unsure these conditions exist.

CONIFERS AND TREES IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS, LANDSCAPES AND PARKS (INCLUDING CHRISTMAS TREES, OAKS and PINES)

APPLICATION GUIDELINES:

Disease Prevention Program*: Apply lower rate at 7-28 day intervals. Do not apply K-PHITE 7LP at intervals less than 3 days.

Disease Control Program: Apply higher rate at 7-14 day intervals until control is reached. Under severe circumstances, application can be made at intervals of up to every three days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity.

Do not apply K-PHITE 7LP at intervals less than 3 days.

K-PHITE 7LP applied in conjunction with good cultural management practices will control Phytophthora root rot of conifers including (but not limited to): Douglas Fir, spruce and pines also other Phytophthora disease of Oaks (Sudden Oak Death) and forest trees. Use the higher rates when disease pressure is severe. Do not apply to trees that are under heat or moisture stress. *Do not graze livestock in treated conifer nurseries and/or plantations. Do not feed forage to livestock from treated nurseries and/or plantations.

DISEASE	APPLICATION METHOD	RATE/APPLICATION
Phytophthora Sudden Oak Death	Foliar Spray	1-2 quarts in a minimum of 100 gallons of water/acre.
	Soil Drench	12-24 fluid ounces/100 gallons of water of 1/8 to 1/4 fluid ounce/gallon of water. Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media.
	Trunk or Bark Injection	3 teaspoons/linear yard of canopy width or 2 inches of trunk diameter at breast height (DBH). Inject directly into tree. Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.
	Root Dip	2 quarts/100 gallons of water or 2/3 fluid ounce/gallon of water. Apply as a pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.

*Do not graze livestock in treated conifer nurseries and/or plantations. Do not feed forage to livestock from treated nurseries and/or plantations.

TURF

K-PHITE 7LP is a systemic Fungicide/Bactericide, which is used for the control of Pythium blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses on sod farms and commercial turf production.

APPLICATION GUIDELINES:

Apply amount in minimum of 2 gallons of water per 1,000 sq. ft.

Disease Prevention Program*: Apply lower rate at 7-28 day intervals. Do not apply K-PHITE 7LP at intervals less than 3 days.

Disease Control Program: Apply higher rate at 7-14 day intervals until control is reached. Under severe circumstances, application can be made at intervals of up to every three days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity.

Do not apply K-PHITE 7LP at intervals less than 3 days.

DISEASE	APPLICATION METHOD	RATE/APPLICATION
Acidovorax Algae Alternaria Anthracnose (Colletotrichum graminicola) Bacterial Wilt Cercospora Helminthosporium Gray Snow Mold Phytophthora Pink Snow Mold	Foliar Spray	3-6 fluid ounces/1000 square feet. (Fairways: 2-6 fluid ounces/1000 square feet.) OR 3-6 quarts/100 gallons of water/acre.
Pythium (Yellow Turf) Rhizoctonia Sclerotinia Xanthomonas Fusarium	Irrigation	Overhead: 2-4 quarts/acre in at least 100 gallons of water.

*Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry place.

PESTICIDE STORAGE: Keep containers tightly closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, keep out of smoke.

NONREFILLABLE CONTAINERS: Do not reuse or refill this container. Clean container promptly after emptying.

NONREFILLABLE CONTAINER EQUAL TO OR LESS THAN 5 GALLONS: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 120 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NONREFILLABLE CONTAINER GREATER THAN 5 GALLONS: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

NOTICE TO BUYER

To the extent permitted by law, all conditions and warranties, and statutory or other rights of action which buyer or any other user may have against Plant Food Systems Inc. or seller are hereby excluded. Plant Food Systems Inc. hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Plant Food Systems Inc. or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Plant Food Systems Inc.'s liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price thereof.